



General Certificate of Education  
Advanced Level Examination  
January 2010

# Mathematics

# MPC4

## Unit Pure Core 4

Tuesday 19 January 2010 9.00 am to 10.30 am

**For this paper you must have:**

- an 8-page answer book
- the blue AQA booklet of formulae and statistical tables.  
You may use a graphics calculator.

**Time allowed**

- 1 hour 30 minutes

**Instructions**

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The **Examining Body** for this paper is AQA. The **Paper Reference** is MPC4.
- Answer **all** questions.
- Show all necessary working; otherwise marks for method may be lost.

**Information**

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

**Advice**

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.

---

Answer **all** questions.

---

1 The polynomial  $f(x)$  is defined by  $f(x) = 15x^3 + 19x^2 - 4$ .

(a) (i) Find  $f(-1)$ . (1 mark)

(ii) Show that  $(5x - 2)$  is a factor of  $f(x)$ . (2 marks)

(b) Simplify

$$\frac{15x^2 - 6x}{f(x)}$$

giving your answer in a fully factorised form. (5 marks)

2 (a) Express  $\cos x + 3 \sin x$  in the form  $R \cos(x - \alpha)$ , where  $R > 0$  and  $0 < \alpha < \frac{\pi}{2}$ . Give your value of  $\alpha$ , in radians, to three decimal places. (3 marks)

(b) (i) Hence write down the minimum value of  $\cos x + 3 \sin x$ . (1 mark)

(ii) Find the value of  $x$  in the interval  $0 \leq x \leq 2\pi$  at which this minimum occurs, giving your answer, in radians, to three decimal places. (2 marks)

(c) Solve the equation  $\cos x + 3 \sin x = 2$  in the interval  $0 \leq x \leq 2\pi$ , giving all solutions, in radians, to three decimal places. (4 marks)

3 (a) (i) Find the binomial expansion of  $(1 + x)^{-\frac{1}{3}}$  up to and including the term in  $x^2$ . (2 marks)

(ii) Hence find the binomial expansion of  $\left(1 + \frac{3}{4}x\right)^{-\frac{1}{3}}$  up to and including the term in  $x^2$ . (2 marks)

(b) Hence show that  $\sqrt[3]{\frac{256}{4 + 3x}} \approx a + bx + cx^2$  for small values of  $x$ , stating the values of the constants  $a$ ,  $b$  and  $c$ . (3 marks)

4 The expression  $\frac{10x^2 + 8}{(x + 1)(5x - 1)}$  can be written in the form  $2 + \frac{A}{x + 1} + \frac{B}{5x - 1}$ , where  $A$  and  $B$  are constants.

(a) Find the values of  $A$  and  $B$ . (4 marks)

(b) Hence find  $\int \frac{10x^2 + 8}{(x + 1)(5x - 1)} dx$ . (4 marks)

5 A curve is defined by the equation

$$x^2 + xy = e^y$$

Find the gradient at the point  $(-1, 0)$  on this curve. (5 marks)

6 (a) (i) Express  $\sin 2\theta$  and  $\cos 2\theta$  in terms of  $\sin \theta$  and  $\cos \theta$ . (2 marks)

(ii) Given that  $0 < \theta < \frac{\pi}{2}$  and  $\cos \theta = \frac{3}{5}$ , show that  $\sin 2\theta = \frac{24}{25}$  and find the value of  $\cos 2\theta$ . (2 marks)

(b) A curve has parametric equations

$$x = 3 \sin 2\theta, \quad y = 4 \cos 2\theta$$

(i) Find  $\frac{dy}{dx}$  in terms of  $\theta$ . (3 marks)

(ii) At the point  $P$  on the curve,  $\cos \theta = \frac{3}{5}$  and  $0 < \theta < \frac{\pi}{2}$ . Find an equation of the tangent to the curve at the point  $P$ . (3 marks)

7 Solve the differential equation  $\frac{dy}{dx} = \frac{1}{y} \cos\left(\frac{x}{3}\right)$ , given that  $y = 1$  when  $x = \frac{\pi}{2}$ .

Write your answer in the form  $y^2 = f(x)$ . (6 marks)

**Turn over for the next question**

**Turn over ►**

- 8 The points  $A$ ,  $B$  and  $C$  have coordinates  $(2, -1, -5)$ ,  $(0, 5, -9)$  and  $(9, 2, 3)$  respectively.

The line  $l$  has equation  $\mathbf{r} = \begin{bmatrix} 2 \\ -1 \\ -5 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ -3 \\ 2 \end{bmatrix}$ .

- (a) Verify that the point  $B$  lies on the line  $l$ . (2 marks)
- (b) Find the vector  $\overrightarrow{BC}$ . (2 marks)
- (c) The point  $D$  is such that  $\overrightarrow{AD} = 2\overrightarrow{BC}$ .
- (i) Show that  $D$  has coordinates  $(20, -7, 19)$ . (2 marks)
- (ii) The point  $P$  lies on  $l$  where  $\lambda = p$ . The line  $PD$  is perpendicular to  $l$ . Find the value of  $p$ . (5 marks)

- 9 A botanist is investigating the rate of growth of a certain species of toadstool. She observes that a particular toadstool of this type has a height of 57 millimetres at a time 12 hours after it begins to grow.

She proposes the model  $h = A\left(1 - e^{-\frac{1}{4}t}\right)$ , where  $A$  is a constant, for the height  $h$  millimetres of the toadstool,  $t$  hours after it begins to grow.

- (a) Use this model to:
- (i) find the height of the toadstool when  $t = 0$ ; (1 mark)
- (ii) show that  $A = 60$ , correct to two significant figures. (2 marks)
- (b) Use the model  $h = 60\left(1 - e^{-\frac{1}{4}t}\right)$  to:
- (i) show that the time  $T$  hours for the toadstool to grow to a height of 48 millimetres is given by
- $$T = a \ln b$$
- where  $a$  and  $b$  are integers; (3 marks)
- (ii) show that  $\frac{dh}{dt} = 15 - \frac{h}{4}$ ; (3 marks)
- (iii) find the height of the toadstool when it is growing at a rate of 13 millimetres per hour. (1 mark)

**END OF QUESTIONS**