



General Certificate of Education
January 2009
Advanced Level Examination

MATHEMATICS
Unit Further Pure 3

MFP3

Wednesday 21 January 2009 1.30 pm to 3.00 pm

For this paper you must have:

- a 12-page answer book
- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 30 minutes

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MFP3.
- Answer **all** questions.
- Show all necessary working; otherwise marks for method may be lost.

Information

- The maximum mark for this paper is 75.
- The marks for questions are shown in brackets.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.

Answer **all** questions.

- 1 The function $y(x)$ satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where

$$f(x, y) = \frac{x^2 + y^2}{x + y}$$

and

$$y(1) = 3$$

- (a) Use the Euler formula

$$y_{r+1} = y_r + hf(x_r, y_r)$$

with $h = 0.2$, to obtain an approximation to $y(1.2)$. (3 marks)

- (b) Use the improved Euler formula

$$y_{r+1} = y_r + \frac{1}{2}(k_1 + k_2)$$


where $k_1 = hf(x_r, y_r)$ and $k_2 = hf(x_r + h, y_r + k_1)$ and $h = 0.2$, to obtain an approximation to $y(1.2)$, giving your answer to four decimal places. (5 marks)

- 2 (a) Show that $\frac{1}{x^2}$ is an integrating factor for the first-order differential equation

$$\frac{dy}{dx} - \frac{2}{x}y = x \quad (3 \text{ marks})$$

- (b) Hence find the general solution of this differential equation, giving your answer in the form $y = f(x)$. (4 marks)

- 3 The diagram shows a sketch of a loop, the pole O and the initial line.



The polar equation of the loop is


$$r = (2 + \cos \theta)\sqrt{\sin \theta}, \quad 0 \leq \theta \leq \pi$$

Find the area enclosed by the loop.

(6 marks)

- 4 (a) Use integration by parts to show that $\int \ln x \, dx = x \ln x - x + c$, where c is an arbitrary constant. (2 marks)
- (b) Hence evaluate $\int_0^1 \ln x \, dx$, showing the limiting process used. (4 marks)

- 5 The diagram shows a sketch of a curve C , the pole O and the initial line.



The curve C has polar equation

$$r = \frac{2}{3 + 2 \cos \theta}, \quad 0 \leq \theta \leq 2\pi$$

- (a) Verify that the point L with polar coordinates $(2, \pi)$ lies on C . (1 mark)
- (b) The circle with polar equation $r = 1$ intersects C at the points M and N .
- (i) Find the polar coordinates of M and N . (3 marks)
- (ii) Find the area of triangle LMN . (4 marks)
- (c) Find a cartesian equation of C , giving your answer in the form $9y^2 = f(x)$. (5 marks)

Turn over for the next question

Turn over ►

6 The function f is defined by $f(x) = e^{2x}(1 + 3x)^{-\frac{2}{3}}$.

(a) (i) Use the series expansion for e^x to write down the first four terms in the series expansion of e^{2x} . (2 marks)

(ii) Use the binomial series expansion of $(1 + 3x)^{-\frac{2}{3}}$ and your answer to part (a)(i) to show that the first three non-zero terms in the series expansion of $f(x)$ are $1 + 3x^2 - 6x^3$. (5 marks)

(b) (i) Given that $y = \ln(1 + 2 \sin x)$, find $\frac{d^2y}{dx^2}$. (4 marks)

(ii) By using Maclaurin's theorem, show that, for small values of x ,

$$\ln(1 + 2 \sin x) \approx 2x - 2x^2 \quad (2 \text{ marks})$$

(c) Find

$$\lim_{x \rightarrow 0} \frac{1 - f(x)}{x \ln(1 + 2 \sin x)} \quad (3 \text{ marks})$$

7 (a) Given that $x = e^t$ and that y is a function of x , show that

$$x^2 \frac{d^2y}{dx^2} = \frac{d^2y}{dt^2} - \frac{dy}{dt} \quad (7 \text{ marks})$$

(b) Hence show that the substitution $x = e^t$ transforms the differential equation

$$x^2 \frac{d^2y}{dx^2} - 4x \frac{dy}{dx} = 10$$

into

$$\frac{d^2y}{dt^2} - 5 \frac{dy}{dt} = 10 \quad (2 \text{ marks})$$

(c) Find the general solution of the differential equation $\frac{d^2y}{dt^2} - 5 \frac{dy}{dt} = 10$. (5 marks)

(d) Hence solve the differential equation $x^2 \frac{d^2y}{dx^2} - 4x \frac{dy}{dx} = 10$, given that $y = 0$ and $\frac{dy}{dx} = 8$ when $x = 1$. (5 marks)

END OF QUESTIONS