

Please write clearly in block capitals.

Centre number

Candidate number

Surname \_\_\_\_\_

Forename(s) \_\_\_\_\_

Candidate signature \_\_\_\_\_

# A-level MATHEMATICS

## Unit Further Pure 3

Wednesday 16 May 2018

Morning

Time allowed: 1 hour 30 minutes

### Materials

For this paper you must have:

- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

### Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

### Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
<b>TOTAL</b>	



Answer **all** questions.

Answer each question in the space provided for that question.

**1 (a)** Explain why  $\int_1^{\infty} \frac{x-1}{e^x} dx$  is an improper integral.

[1 mark]

**(b)** Evaluate the improper integral  $\int_1^{\infty} \frac{x-1}{e^x} dx$ , showing the limiting process used.

[4 marks]

QUESTION  
PART  
REFERENCE

**Answer space for question 1**





2 It is given that  $y(x)$  satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where  $f(x, y) = x + \frac{1}{2} \log_2(y + 7)$

and  $y(2) = 1$

Use the improved Euler formula

$$y_{r+1} = y_r + \frac{1}{2}(k_1 + k_2)$$


where  $k_1 = hf(x_r, y_r)$  and  $k_2 = hf(x_r + h, y_r + k_1)$  and  $h = 0.3$ , to obtain an approximation to  $y(2.3)$ , giving your answer to four significant figures.

[5 marks]

QUESTION  
PART  
REFERENCE

Answer space for question 2





QUESTION  
PART  
REFERENCE

**Answer space for question 2**

Answer space consisting of 25 horizontal lines for writing.

**Turn over ▶**



3 Find the general solution of the differential equation

$$\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 10y = 34x - 20x^2$$

[7 marks]

QUESTION  
PART  
REFERENCE

Answer space for question 3





4 Show that, for some value of  $k$ ,

$$\lim_{x \rightarrow 0} \left[ \frac{3 - \sqrt{9 - kx^4}}{7x^6 + 8x^4} \right] = \frac{1}{32}$$

and state this value of  $k$ .

[4 marks]

QUESTION  
PART  
REFERENCE

Answer space for question 4







5 The polar equation of an ellipse  $C$  is

$$r = \frac{5}{3 + 2 \sin \theta}, \quad 0 \leq \theta \leq 2\pi$$

(a) By finding a Cartesian equation of the ellipse  $C$  in a suitable form, state the Cartesian equations of the tangents to  $C$  that are parallel to the coordinate axes. [5 marks]

(b) Given that the area of the region bounded by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is  $\pi ab$ , deduce the exact value of  $\int_0^{2\pi} \frac{1}{(3 + 2 \sin \theta)^2} d\theta$ . [4 marks]

QUESTION  
PART  
REFERENCE

**Answer space for question 5**





6 A second order differential equation is given by

$$\frac{d^2y}{dx^2} + (\cot x + \tan x) \frac{dy}{dx} = y \cot^2 x, \quad 0 < x < \frac{\pi}{2}$$

(a) Show that the substitution

$$u = \frac{dy}{dx} + y \cot x$$

transforms the second order differential equation into

$$\frac{du}{dx} + u \tan x = 0$$

[3 marks]

(b) Hence, given that  $y = 0$  and  $\frac{dy}{dx} = \sqrt{3}$  when  $x = \frac{\pi}{6}$ , solve the second order differential equation to find an expression for  $y$  in terms of  $\sin x$ .

[9 marks]

QUESTION  
PART  
REFERENCE

Answer space for question 6



QUESTION  
PART  
REFERENCE

Answer space for question 6


Turn over ►







7 (a) Find the general solution of the differential equation

$$\frac{d^2y}{dx^2} + 4 \frac{dy}{dx} + 4y = 4 \sin 2x + 8 \cos 2x$$

[6 marks]

(b) It is given that  $y = f(x)$  is the solution of the differential equation

$$\frac{d^2y}{dx^2} + 4 \frac{dy}{dx} + 4y = 4 \sin 2x + 8 \cos 2x$$

such that the first two non-zero terms in the Maclaurin's series, in ascending powers of  $x$ , of  $f(x)$  are  $\frac{1}{2} + kx^2$ . Find the value of  $f\left(\frac{\pi}{6}\right)$ , giving your answer in an exact form.

[4 marks]

QUESTION  
PART  
REFERENCE


Answer space for question 7







- 8 The diagram shows the sketch of a curve  $C_1$ .



The polar equation of the curve  $C_1$  is  $r = 4 \cos^2 \theta$ ,  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ .

The curve  $C_2$  has polar equation  $r = 1 + \tan \theta$ ,  $0 \leq \theta < \frac{\pi}{2}$ .

- (a) Prove that the curves  $C_1$  and  $C_2$  intersect at a single point  $P$  whose distance from the pole  $O$  is 2. [5 marks]
- (b) The curve  $C_2$  meets the initial line at the point  $A$ . Find the area of the region bounded by the line segment  $AP$  and the curve  $C_2$ , giving your answer in an exact form. [8 marks]

QUESTION  
PART  
REFERENCE

**Answer space for question 8**



QUESTION PART REFERENCE	Answer space for question 8

**Turn over ►**



QUESTION  
PART  
REFERENCE

### Answer space for question 8

Turn over ►



2 1

- 9 (a)** Find the first three non-zero terms in the expansion of

$$\ln(1 + y) - \ln(1 - y)$$

in ascending powers of  $y$  **and** state the range of values of  $y$  for which the expansion is valid.

[3 marks]

- (b)** Use the identity  $1 + x^3 \equiv (1 + x)(1 - x + x^2)$  to find the coefficient of  $x^{6r-3}$ , where  $r$  is a positive integer, in the expansion of

$$\ln\left(\frac{1 - x + x^2}{1 + x + x^2}\right)$$

in ascending powers of  $x$ . Give your answer in its simplest form.

[7 marks]

QUESTION  
PART  
REFERENCE

**Answer space for question 9**





