



Pearson
Edexcel

Mark Scheme (Results)

Summer 2018

Pearson Edexcel GCE Further Mathematics
AS Further Mechanics M2 Paper 8FM0_26

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2018

Publications Code 8FM0_26_1806_MS

All the material in this publication is copyright

© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the last candidate in exactly the same way as they mark the first.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification/indicative content will not be exhaustive.

General Instructions for Marking

1. The total number of marks for the paper is 40.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.


- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 5. Where a candidate has made multiple responses and indicates which response they wish to submit, examiners should mark this response. If there are several attempts at a question which have not been crossed out, examiners should mark the final answer which is the answer that is the most complete.
 6. Ignore wrong working or incorrect statements following a correct answer.
 7. Mark schemes will firstly show the solution judged to be the most common response expected from candidates. Where appropriate, alternatives answers are provided in the notes. If examiners are not sure if an answer is acceptable, they will check the mark scheme to see if an alternative answer is given for the method used.

Qu	Scheme	Marks	AOs	Notes
1(a)	Complete strategy to find d	M1	3.1b	Complete strategy to find d e.g. moments about AB or a parallel axis. Needs all relevant terms. Must be dimensionally correct. Condone sign errors. M 's might cancel from the start.
	$\frac{5}{30}M \times \frac{5}{2}a + \frac{13}{30}M \times \frac{5}{2}a = M \times d$	A1	1.1b	Unsimplified equation with at most one error
	$\left(\frac{25}{2}a + \frac{65}{2}a = 30d\right)$	A1	1.1b	Correct unsimplified equation
	$90a = 60d \Rightarrow d = \frac{3}{2}a$ *	A1*	2.1	Obtain the given answer from a convincing argument
		(4)		
1(b)	Complete strategy to find k , e.g. by use of a moments equation	M1	3.1b	Complete strategy to find k e.g. moments about A . Needs all relevant terms. Must be dimensionally correct. Condone sign errors. Condone if a, M, g missing throughout
	$Mg \times \frac{3}{2}a = kMg \times 12a$	A1	1.1b	Correct unsimplified equation in k
	$k = \frac{1}{8}$	A1	1.1b	Correct answer – any equivalent form
		(3)		
1(b) alt	Moments equation	M1		
	$12a \times kM = \frac{13}{30}M \times 2.5a + \frac{5}{30}M \times 2.5a$	A1		
	$12k = \frac{45}{30}, k = \frac{1}{8}$	A1		
		(3)		
			(7 marks)	

Qu	Scheme	Marks	AOs	Notes
2(a)	Complete strategy to find value of θ	M1	3.1b	Complete strategy involving resolving in perpendicular directions, change of units and solution of simultaneous equations
	Resolve vertically	M1	3.1b	Complete strategy to form one equation involving θ e.g. resolve vertically. Condone sin/cos confusion
	$R \cos \theta^\circ = mg$	A1	1.1b	Or equivalent
	Resolve horizontally	M1	3.1b	Complete strategy to form a second equation involving θ e.g. resolve horizontally. Condone sin/cos confusion
	$R \sin \theta^\circ = \frac{mv^2}{r}$	A1	1.1b	Correct unsimplified – need not substitute for v or r
	$v = 80 \text{ km h}^{-1} = \frac{80 \times 1000}{60^2} \text{ m s}^{-1}$	B1	1.2	Correct conversion km h^{-1} to m s^{-1} (22.2)
	Solve simultaneous equations and substitute v in correct units to obtain θ : $\tan \theta^\circ = \frac{v^2}{rg} = \frac{640000}{36^2 \times 500 \times 9.8}$, $\theta = 5.8$	A1	2.2a	Accept 5.8 or 5.75 (follows use of 9.8)
		(7)		
(b)	All weight acting at a single point	B1	3.5b	Any appropriate comment e.g. Only one point of contact with the road The centre of mass of the car is on the road.
		(1)		
(c)	Friction acting down the slope	B1	2.2a	Need to include the direction
		(1)		
(9 marks)				

Qu	Scheme				Marks	AOs	Notes
3(a)	L is symmetrical about AD				B1	2.4	Any equivalent statement about the symmetry
					(1)		
3(b)		$ABDF$	BCD	DEF	L		
	Mass ratio	$4a^2 \times M$	$a^2 \times 3M$	$a^2 \times 3M$	$10a^2 \times M$		
	C of M from BE	$-a$	$+\frac{a}{3}$	$-\frac{2a}{3}$	x		
	Mass ratios				B1	1.2	Correct mass ratios
	Distances from BE				B1	1.2	Distance ratios from any horizontal or vertical axis
	Moments equation				M1	2.1	Moments equation for complete lamina about any horizontal or vertical axis. Must be dimensionally correct
	$-a \times 4a^2 M + \frac{a}{3} \times 3a^2 M - \frac{2a}{3} \times 3a^2 M = 10a^2 M \times x$ $(-4a + a - 2a = 10x)$				A1	1.1b	Correct unsimplified equation for their axes
	$x = -\frac{5a}{10} = -\frac{a}{2}$				A1	1.1b	Correct horizontal or vertical distance from D
	Use symmetry and Pythagoras				M1	1.1a	Use of Pythagoras with their distance
Distance from $D = \sqrt{\frac{a^2}{4} + \frac{a^2}{4}} = \frac{\sqrt{2}}{2} a$ *				A1*	2.2a	Obtain given answer from correct working.	
				(7)			

3(c)



Trig ratio of a relevant angle	M1	1.2	Trig ratio of θ or $90^\circ - \theta$ or equivalent
$\tan \theta = \frac{1}{3} \quad \text{or} \quad \cos \theta = \frac{\frac{10}{4}a^2 + 4a^2 - \frac{2}{4}a^2}{2 \times \frac{\sqrt{10}}{2}a \times 2a} = \frac{6}{2\sqrt{10}}$	A1ft	1.1b	Correct unsimplified expression using their $\frac{a}{2}$
$\theta = 18.4^\circ$	A1	1.1b	Correct angle. Accept 0.322 radians
	(3)		

(11 marks)

Qu	Scheme	Marks	AOs	Notes
4(a)	$\int \frac{1}{4} dt = \int \frac{1}{50-10v} dv$	M1	3.1a	Strategy to find v and attempt the integration
	$\frac{1}{4}t = -\frac{1}{10} \ln(50-10v) (+C)$	A1	1.1b	Correct integration
	$\frac{1}{4}t = -\frac{1}{10} \ln(50-10v) + \frac{1}{10} \ln 50$	M1	1.1b	Use boundary conditions as limits or evaluate constant of integration in an expression involving $\lambda \ln(a+bv)$ and μt
	$-\frac{5t}{2} = \ln\left(\frac{5-v}{5}\right)$	M1	1.1b	Remove logarithm to express v in terms of t
	$v = 5(1 - e^{-2.5t})$ *	A1*	2.1	Obtain given answer from correct working
		(5)		
4(b)	limiting value is 5	B1	2.2a	Correct answer from correct working
		(1)		
4(c)	Equation in x and t : $\frac{dx}{dt} = 5(1 - e^{-2.5t})$	M1	1.1a	Set up equation of motion in terms of x and t
	$\Rightarrow \int 1 dx = \int 5(1 - e^{-2.5t}) dt$	M1	1.1b	Separate variables and attempt integration of both sides
	$x = 5t + 2e^{-2.5t} (+C)$	A1	1.1b	Any equivalent form. Condone if $+C$ not seen
	Use $v = 2.5$ and $v = 5(1 - e^{-2.5t})$ to find value of t	M1	3.1a	Use $v = 2.5$ to find limit for t
	$1 - \frac{2.5}{5} = e^{-2.5t} \Rightarrow t = \frac{2}{5} \ln 2$	A1	1.1b	Any equivalent exact form. (0.277)
	$[x]_0^d = [5t + 2e^{-2.5t}]_0^{\frac{2}{5} \ln 2}$	M1	2.1	Use boundary conditions as limits or evaluate constant of integration in an expression involving λt and $\mu e^{-2.5t}$
	$d = 2 \ln 2 - 1$ *	A1*	1.1b	Sufficient correct working to justify given answer
		(7)		
(13 marks)				

