

Mark Scheme (Results)

June 2014

Pearson Edexcel GCE in Core Mathematics 4R
(6666/01R)

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2014

Publications Code UA038470

All the material in this publication is copyright

© Pearson Education Ltd 2014

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

SOME GENERAL PRINCIPLES FOR CORE MATHS MARKING

(But the particular mark scheme always takes precedence)

Method mark for solving 3 term quadratic:

Question Number	Scheme
3(b)	<p>Factorising/Solving a quadratic equation is tested in Question 3(b).</p> <p>Method mark for solving a 3 term quadratic:</p> <p>1. Factorisation $(x^2 + bx + c) = (x + p)(x + q)$, where $pq = c$, leading to $x = \dots$ $(ax^2 + bx + c) = (mx \pm p)(nx \pm q)$, where $pq = c$ and $mn = a$, leading to $x = \dots$</p> <p>2. Formula Attempt to use correct formula (with values for a, b and c.)</p> <p>3. Completing the square Solving $x^2 + bx + c = 0$: $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$, $q \neq 0$, leading to $x = \dots$</p>

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are mistakes in the substitution of values (but refer to the mark scheme first... the application of this principle may vary).

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but will be lost if there is any mistake in the working.

Answers without working

The rubric says that these may gain no credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice.

Misreads

A misread must be consistent for the whole question to be interpreted as such.

These are not common. In clear cases, please deduct the first 2 A (or B) marks which would have been lost by following the scheme. (Note that 2 marks is the maximum misread penalty, but that misreads which alter the nature or difficulty of the question cannot be treated so generously and it will usually be necessary here to follow the scheme as written).

Sometimes following the scheme as written is more generous to the candidate than applying the misread rule, so in this case use the scheme as written.

Question Number	Scheme	Marks
<p>1. (a)</p> <p>(b)</p>	$\left\{ \frac{1}{\sqrt{(9-10x)}} \right\} (9-10x)^{-\frac{1}{2}}$ $= (9)^{-\frac{1}{2}} \left(1 - \frac{10x}{9} \right)^{-\frac{1}{2}} = \frac{1}{3} \left(1 - \frac{10x}{9} \right)^{-\frac{1}{2}}$ $= \left\{ \frac{1}{3} \right\} \left[1 + \left(-\frac{1}{2} \right) (kx) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!} (kx)^2 + \dots \right]$ $= \left\{ \frac{1}{3} \right\} \left[1 + \left(-\frac{1}{2} \right) \left(\frac{-10x}{9} \right) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!} \left(\frac{-10x}{9} \right)^2 + \dots \right]$ $= \frac{1}{3} \left[1 + \frac{5}{9}x + \frac{25}{54}x^2 + \dots \right]$ $= \frac{1}{3} + \frac{5}{27}x + \frac{25}{162}x^2 + \dots$ $\frac{3+x}{\sqrt{(9-10x)}} = (3+x)(9-10x)^{-\frac{1}{2}}$ $= (3+x) \left(\frac{1}{3} + \frac{5}{27}x + \left\{ \frac{25}{162}x^2 + \dots \right\} \right)$ $= 1 + \frac{5}{9}x + \frac{25}{54}x^2 + \frac{1}{3}x + \frac{5}{27}x^2 + \dots$ $= 1 + \frac{8}{9}x + \frac{35}{54}x^2 + \dots$	<p>$(9-10x)^{-\frac{1}{2}}$ or uses power of $-\frac{1}{2}$ B1</p> <p>$(9)^{-\frac{1}{2}}$ or $\frac{1}{3}$ B1</p> <p>At least two correct terms. See notes M1</p> <p>A1; A1 [5]</p> <p><i>Can be implied by later work</i> See notes M1</p> <p>Multiplies out to give exactly one constant term, exactly 2 terms in x and exactly 2 terms in x^2. M1</p> <p>Ignore terms in x^3. Can be implied. A1</p> <p>[3] 8</p>
Question 1 Notes		
(a)	<p>B1 Writes down $(9-10x)^{-\frac{1}{2}}$ or uses power of $-\frac{1}{2}$.</p> <p>This mark can be implied by a constant term of $(9)^{-\frac{1}{2}}$ or $\frac{1}{3}$.</p> <p>B1 $(9)^{-\frac{1}{2}}$ or $\frac{1}{3}$ outside brackets or $\frac{1}{3}$ as candidate's constant term in their binomial expansion.</p> <p>M1 Expands $(\dots + kx)^{-\frac{1}{2}}$ to give any 2 terms out of 3 terms simplified or an un-simplified, $1 + (-\frac{1}{2})(kx)$ or $(-\frac{1}{2})(kx) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!}(kx)^2$ or $1 + \dots + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!}(kx)^2$, where $k \neq 1$.</p> <p>A1 $\frac{1}{3} + \frac{5}{27}x$ (simplified fractions)</p> <p>A1 Accept only $\frac{25}{162}x^2$</p>	

1. (a) ctd	<p>Note You cannot recover correct work for part (a) in part (b). i.e. if the correct answer to (a) appears as part of their solution in part (b), it cannot be credited in part (a).</p> <p>SC If a candidate <i>would otherwise score</i> A0A0 then allow Special Case 1st A1 for either</p> <p>SC: $\frac{1}{3}\left[1 + \frac{5}{9}x; \dots\right]$ or SC: $\lambda\left[1 + \frac{5}{9}x + \frac{25}{54}x^2 + \dots\right]$ or SC: $\left[\lambda + \frac{5\lambda}{9}x + \frac{25\lambda}{54}x^2 + \dots\right]$</p> <p>(where λ can be 1 or omitted), with each term in the [.....] is a simplified fraction</p> <p>SC Special case for the M1 mark</p> <p>Award Special Case M1 for a correct simplified or un-simplified $1 + n(kx) + \frac{n(n-1)}{2!}(kx)^2$ expansion with a value of $n \neq -\frac{1}{2}$, $n \neq$ positive integer and a consistent (kx). Note that (kx) must be consistent (on the RHS, not necessarily the LHS) in a candidate's expansion. Note that $k \neq 1$.</p> <p>Note Candidates who write $\left\{\frac{1}{3}\right\}\left[1 + \left(-\frac{1}{2}\right)\left(\frac{10x}{9}\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}\left(\frac{10x}{9}\right)^2 + \dots\right]$ where $k = \frac{10}{9}$ and not $-\frac{10}{9}$ and achieve $\frac{1}{3} - \frac{5}{27}x; + \frac{25}{162}x^2 + \dots$ will get B1B1M1A0A1.</p>
(b)	<p>M1 Writes down $(3 + x)$(their part (a) answer, at least 2 of the 3 terms.)</p> <p>Note $(3 + x)\left(\frac{1}{4} + \frac{5}{4}x + \dots\right)$ or $(3 + x)\left(\frac{1}{3} + \frac{5}{27}x + \frac{25}{162}x^2 + \dots\right)$ are fine for M1.</p> <p>Note This mark can also be implied by candidate multiplying out to find two terms (or coefficients) in x.</p> <p>M1 Multiplies out to give exactly one constant term, exactly 2 terms in x and exactly 2 terms in x^2.</p> <p>Note This M1 mark can be implied. You can also ignore x^3 terms.</p> <p>A1 $1 + \frac{8}{9}x + \frac{35}{54}x^2 + \dots$</p> <p>Alternative Methods for part (a) Alternative method 1: Candidates can apply an alternative form of the binomial expansion.</p> $\left\{\frac{1}{\sqrt{(9-10x)}}\right\} = (9-10x)^{-\frac{1}{2}} = (9)^{-\frac{1}{2}} + \left(-\frac{1}{2}\right)(9)^{-\frac{3}{2}}(-10x) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}(9)^{-\frac{5}{2}}(-10x)^2$ <p>B1 Writes down $(9-10x)^{-\frac{1}{2}}$ or uses power of $-\frac{1}{2}$.</p> <p>B1 $9^{-\frac{1}{2}}$ or $\frac{1}{3}$</p> <p>M1 Any two of three (un-simplified or simplified) terms correct.</p> <p>A1 $\frac{1}{3} + \frac{5}{27}x$</p> <p>A1 $\frac{25}{162}x^2$</p> <p>Note The terms in C need to be evaluated, so $^{-\frac{1}{2}}C_0(9)^{-\frac{1}{2}} + ^{-\frac{1}{2}}C_1(9)^{-\frac{3}{2}}(-10x) + ^{-\frac{1}{2}}C_2(9)^{-\frac{5}{2}}(-10x)^2$ without further working is B1B0M0A0A0.</p>

<p>1. (a)</p>	<p>Alternative Method 2: Maclaurin Expansion</p> <p>Let $f(x) = \frac{1}{\sqrt{(9-10x)}}$</p> <p>$\{f(x) = (9-10x)^{-\frac{1}{2}}\}$</p> <p>$f''(x) = 75(9-10x)^{-\frac{5}{2}}$</p> <p>$f'(x) = (-\frac{1}{2})(9-10x)^{-\frac{3}{2}}(-10)$</p> <p>$\left\{ \therefore f(0) = \frac{1}{3}, f'(0) = \frac{5}{27} \text{ and } f''(0) = \frac{75}{243} = \frac{25}{81} \right\}$</p> <p>$f(x) = \frac{1}{3} + \frac{5}{27}x + \frac{25}{162}x^2 + \dots$</p>	<p>$(9-10x)^{-\frac{1}{2}}$ B1</p> <p>Correct $f''(x)$ B1 oe</p> <p>$\pm a(9-10x)^{-\frac{3}{2}}; a \neq \pm 1$ M1</p> <p>A1; A1</p>
---------------	--	---

Question Number	Scheme	Mark
2. (a)	$\text{Area} \approx \frac{1}{2} \times 0.5 \times [2 + 2(4.077 + 7.389 + 10.043) + 0]$ $= \frac{1}{4} \times 45.018 = 11.2545 = 11.25 \text{ (2 dp)}$	B1; M1 A1 cao [3]
(b)	Any one of <ul style="list-style-type: none"> • Increase the number of strips • Use more trapezia • Make h smaller • Increase the number of x and/or y values used • Shorter /smaller intervals for x • More values of y. • More intervals of x • Increase n 	B1 [1]
(c)	$\left\{ \int (2-x)e^{2x} dx \right\}, \left\{ \begin{array}{l} u = 2-x \Rightarrow \frac{du}{dx} = -1 \\ \frac{dv}{dx} = e^{2x} \Rightarrow v = \frac{1}{2}e^{2x} \end{array} \right\}$	M1 A1 A1 oe dM1 A1 oe [5] 9
Either $(2-x)e^{2x} \rightarrow \pm \lambda(2-x)e^{2x} \pm \int \mu e^{2x} \{dx\}$ or $\pm x e^{2x} \rightarrow \pm \lambda x e^{2x} \pm \int \mu e^{2x} \{dx\}$		
$= \frac{1}{2}(2-x)e^{2x} - \int -\frac{1}{2}e^{2x} \{dx\}$		
$(2-x)e^{2x} \rightarrow \frac{1}{2}(2-x)e^{2x} - \int -\frac{1}{2}e^{2x} \{dx\}$		
$= \frac{1}{2}(2-x)e^{2x} + \frac{1}{4}e^{2x}$		
$\text{Area} = \left\{ \left[\frac{1}{2}(2-x)e^{2x} + \frac{1}{4}e^{2x} \right]_0^2 \right\}$		
$= \left(0 + \frac{1}{4}e^4 \right) - \left(\frac{1}{2}(2)e^0 + \frac{1}{4}e^0 \right)$		
$= \frac{1}{4}e^4 - \frac{5}{4}$		

Question 2 Notes

(a)	B1 M1 Note A1 Note Note	Outside brackets $\frac{1}{2} \times 0.5$ or $\frac{0.5}{2}$ or 0.25 or $\frac{1}{4}$. For structure of trapezium rule [.....]. Condone missing 0. No errors are allowed [eg. an omission of a y-ordinate or an extra y-ordinate or a repeated y ordinate]. 11.25 cao Working must be seen to demonstrate the use of the trapezium rule. The actual area is 12.39953751... Award B1M1A1 for $\frac{0.5}{2}(2+0) + \frac{1}{2}(4.077 + 7.389 + 10.043) = 11.25$
-----	--	---

2. (a) contd	<p>Bracketing mistake: Unless the final answer implies that the calculation has been done correctly. Award B1M0A0 for $\frac{1}{2} \times 0.5 + 2 + 2(4.077 + 7.389 + 10.043) + 0$ (nb: answer of 45.268).</p>
	<p>Alternative method for part (a): Adding individual trapezia</p> $\text{Area} \approx 0.5 \times \left[\frac{2+4.077}{2} + \frac{4.077+7.389}{2} + \frac{7.389+10.043}{2} + \frac{10.043+0}{2} \right] = 11.2545 = 11.25 \text{ (2 dp) cao}$ <p>B1 0.5 and a divisor of 2 on all terms inside brackets. M1 First and last ordinates once and the middle ordinates twice inside brackets ignoring the 2. A1 11.25 cao</p>
(b)	<p>B0 Give B0 for</p> <ul style="list-style-type: none"> • smaller values of x and/or y. • use more decimal places
(c)	<p>M1 Either $(2-x)e^{2x} \rightarrow \pm \lambda(2-x)e^{2x} \pm \int \mu e^{2x} \{dx\}$ or $\pm xe^{2x} \rightarrow \pm \lambda xe^{2x} \pm \int \mu e^{2x} \{dx\}$</p> <p>A1 $(2-x)e^{2x} \rightarrow \frac{1}{2}(2-x)e^{2x} - \int -\frac{1}{2}e^{2x} \{dx\}$ either un-simplified or simplified.</p> <p>A1 Correct expression, i.e. $\frac{1}{2}(2-x)e^{2x} + \frac{1}{4}e^{2x}$ or $\frac{5}{4}e^{2x} - xe^{2x}$ (or equivalent)</p> <p>dM1 which is dependent on the 1st M1 mark being awarded. Complete method of applying limits of 2 and 0 to all terms and subtracting the correct way round.</p> <p>Note Evidence of a proper consideration of the limit of 0 is needed for M1. So, just subtracting zero is M0.</p> <p>A1 $\frac{1}{4}e^4 - \frac{5}{4}$ or $\frac{e^4 - 5}{4}$. Do not allow $\frac{1}{4}e^4 - \frac{5}{4}e^0$ unless simplified to give $\frac{1}{4}e^4 - \frac{5}{4}$</p> <p>Note 12.39953751... without seeing $\frac{1}{4}e^4 - \frac{5}{4}$ is A0.</p> <p>Note 12.39953751... from NO working is M0A0A0M0A0.</p>

Question Number	Scheme	Marks
<p>3.</p> <p>(a)</p> <p>(b)</p>	$x^2 + y^2 + 10x + 2y - 4xy = 10$ $\left\{ \begin{array}{l} \cancel{dx} \\ \cancel{dx} \end{array} \right\} \times \left\{ \begin{array}{l} 2x + 2y \frac{dy}{dx} + 10 + 2 \frac{dy}{dx} - \left(4y + 4x \frac{dy}{dx} \right) = 0 \\ 2x + 10 - 4y + (2y + 2 - 4x) \frac{dy}{dx} = 0 \\ \frac{dy}{dx} = \frac{2x + 10 - 4y}{4x - 2y - 2} \\ \text{Simplifying gives } \frac{dy}{dx} = \frac{x + 5 - 2y}{2x - y - 1} \left\{ = \frac{-x - 5 + 2y}{-2x + y + 1} \right\}$ $\left\{ \frac{dy}{dx} = 0 \Rightarrow \right\} x + 5 - 2y = 0$ <p>So $x = 2y - 5$,</p> $(2y - 5)^2 + y^2 + 10(2y - 5) + 2y - 4(2y - 5)y = 10$ $4y^2 - 20y + 25 + y^2 + 20y - 50 + 2y - 8y^2 + 20y = 10$ <p>gives $-3y^2 + 22y - 35 = 0$ or $3y^2 - 22y + 35 = 0$</p> $(3y - 7)(y - 5) = 0 \text{ and } y = \dots$ $y = \frac{7}{3}, 5$	<p>See notes</p> <p>M1 <u>A1</u> <u>M1</u></p> <p>Dependent on the first M1 mark.</p> <p>dM1</p> <p>A1 cso oe</p> <p>[5]</p> <p>M1</p> <p>M1</p> <p>$3y^2 - 22y + 35 \{= 0\}$</p> <p>A1 oe</p> <p>see notes</p> <p>Method mark for solving a quadratic equation.</p> <p>ddM1</p> <p>$\{y = \} \frac{7}{3}, 5$</p> <p>A1 cao</p> <p>[5]</p>
<p>(b)</p>	<p>Alternative method for part (b)</p> $\left\{ \frac{dy}{dx} = 0 \Rightarrow \right\} x + 5 - 2y = 0$ <p>So $y = \frac{x + 5}{2}$,</p> $x^2 + \left(\frac{x + 5}{2} \right)^2 + 10x + 2 \left(\frac{x + 5}{2} \right) - 4x \left(\frac{x + 5}{2} \right) = 10$ $x^2 + \frac{x^2 + 10x + 25}{4} + 10x + x + 5 - 2x^2 - 10x = 10$ $4x^2 + x^2 + 10x + 25 + 40x + 4x + 20 - 8x^2 - 40x = 40$ <p>gives $-3x^2 + 14x + 5 = 0$ or $3x^2 - 14x - 5 = 0$</p> $(3x + 1)(x - 5) = 0, x = \dots$ $y = \frac{-\frac{1}{3} + 5}{2}, \frac{5 + 5}{2}$ $y = \frac{7}{3}, 5$	<p>M1</p> <p>M1</p> <p>$3x^2 - 14x - 5 \{= 0\}$</p> <p>A1 oe</p> <p>see notes</p> <p>Solves a quadratic and finds at least one value for y.</p> <p>ddM1</p> <p>$\{y = \} \frac{7}{3}, 5$</p> <p>A1 cao</p> <p>[5]</p>
		<p>10</p>

Question 3 Notes	
3. (a)	<p>M1 Differentiates implicitly to include either $\pm 4x \frac{dy}{dx}$ or $y^2 \rightarrow 2y \frac{dy}{dx}$ or $2y \rightarrow 2 \frac{dy}{dx}$. (Ignore $\left(\frac{dy}{dx} = \right)$).</p> <p>A1 $x^2 + y^2 + 10x + 2y \rightarrow 2x + 2y \frac{dy}{dx} + 10 + 2 \frac{dy}{dx}$ and $10 \rightarrow 0$</p> <p>M1 $-4xy \rightarrow \pm 4y \pm 4x \frac{dy}{dx}$</p> <p>Note If an extra term appears then award 1st A0.</p> <p>Note $2x + 2y \frac{dy}{dx} + 10 + 2 \frac{dy}{dx} - 4y - 4x \frac{dy}{dx} \rightarrow 2x + 10 - 4y = -2y \frac{dy}{dx} - 2 \frac{dy}{dx} + 4x \frac{dy}{dx}$ will get 1st A1 (implied) as the "= 0" can be implied by rearrangement of their equation.</p> <p>dm1 dependent on the first method mark being awarded. An attempt to factorise out all the terms in $\frac{dy}{dx}$ as long as there are at least two terms in $\frac{dy}{dx}$.</p> <p>A1 $\frac{x + 5 - 2y}{2x - y - 1}$ or $\frac{-x - 5 + 2y}{-2x + y + 1}$ (must be simplified).</p> <p>cso: If the candidate's solution is not completely correct, then do not give this mark.</p>
(b)	<p>M1 Sets the numerator of their $\frac{dy}{dx}$ equal to zero (or the denominator of their $\frac{dx}{dy}$ equal to zero) oe.</p> <p>NOTE If the numerator involves one variable only then only the 1st M1 mark is possible in part (b).</p> <p>M1 Substitutes their x or their y into the printed equation to give an equation in one variable only.</p> <p>A1 For obtaining either $-3y^2 + 22y - 35 \{=0\}$ or $3y^2 - 22y + 35 \{=0\}$</p> <p>Note This mark can also awarded for a correct three term equation, eg. either $-3y^2 + 22y = 35$ $3y^2 - 22y = -35$ or $3y^2 + 35 = 22y$ are all fine for A1.</p> <p>ddM1 Dependent on the previous 2 M marks. See notes at the beginning of the mark scheme: Method mark for solving a 3 term quadratic</p> <ul style="list-style-type: none"> • $(3y - 7)(y - 5) = 0 \Rightarrow y = \dots$ • $y = \frac{22 \pm \sqrt{(-22)^2 - 4(3)(35)}}{2(3)}$ • $y^2 - \frac{22}{3}y - \frac{35}{3} = 0 \Rightarrow \left(y - \frac{11}{3}\right)^2 - \frac{121}{9} + \frac{35}{3} = 0 \Rightarrow y = \dots$ • Or writes down at least one correct y-root from their quadratic equation. This is usually found from their calculator. <p>Note If a candidate applies the alternative method then they also need to use their $y = \frac{x + 5}{2}$ in order to find at least one value for y in order to gain the final M1.</p> <p>A1 $y = \frac{7}{3}, 5$. cao. (2.33 or 2.3 without reference to $\frac{7}{3}$ or $2\frac{1}{3}$ is not allowed for this mark.)</p> <p>Note It is possible for a candidate who does not achieve full marks in part (a), (but has a correct numerator for $\frac{dy}{dx}$) to gain all 5 marks in part (b).</p>

Question Number	Scheme	Mark
4. (a)	$\frac{25}{x^2(2x+1)} \equiv \frac{A}{x} + \frac{B}{x^2} + \frac{C}{(2x+1)}$ $B = 25, C = 100$	At least one of "B" or "C" correct. B1 Breaks up their partial fraction correctly into three terms and both "B" = 25 and "C" = 100. B1 cso See notes.
	$25 \equiv Ax(2x+1) + B(2x+1) + Cx^2$ $x=0, \quad 25 = B$ $x = -\frac{1}{2}, \quad 25 = \frac{1}{4}C \Rightarrow C = 100$ $x^2 \text{ terms: } 0 = 2A + C$ $0 = 2A + 100 \Rightarrow A = -50$ $x^2: 0 = 2A + C, \quad x: 0 = A + 2B,$ $\text{constant: } 25 = B$	Writes down a correct identity and attempts to find the value of either one of "A", "B" or "C". M1
	leading to $A = -50$ $\left\{ \frac{25}{x^2(2x+1)} \equiv -\frac{50}{x} + \frac{25}{x^2} + \frac{100}{(2x+1)} \right\}$	Correct value for "A" which is found using a correct identity and follows from their partial fraction decomposition. A1 [4]
(b)	$V = \pi \int_1^4 \left(\frac{5}{x\sqrt{2x+1}} \right)^2 dx$ <p>For their partial fraction</p>	For $\pi \int \left(\frac{5}{x\sqrt{2x+1}} \right)^2 dx$ B1 Ignore limits and dx. Can be implied.
	$\left\{ \int \frac{25}{x^2(2x+1)} dx = \int -\frac{50}{x} + \frac{25}{x^2} + \frac{100}{(2x+1)} dx \right\}$ $= -50 \ln x + \frac{25x^{-1}}{(-1)} + \frac{100}{2} \ln(2x+1) \{+ c\}$	Either $\pm \frac{A}{x} \rightarrow \pm a \ln x$ or $\pm a \ln kx$ or $\pm \frac{B}{x^2} \rightarrow \pm b x^{-1}$ or $\frac{C}{(2x+1)} \rightarrow \pm c \ln(2x+1)$ M1 At least two terms correctly integrated A1ft All three terms correctly integrated. A1ft
	$\left\{ \int_1^4 \frac{25}{x^2(2x+1)} dx = \left[-50 \ln x - \frac{25}{x} + 50 \ln(2x+1) \right]_1^4 \right\}$ $= \left(-50 \ln 4 - \frac{25}{4} + 50 \ln 9 \right) - \left(0 - 25 + 50 \ln 3 \right)$ $= 50 \ln 9 - 50 \ln 4 - 50 \ln 3 - \frac{25}{4} + 25$ $= 50 \ln \left(\frac{3}{4} \right) + \frac{75}{4}$ $\text{So, } V = \frac{75}{4} \pi + 50 \pi \ln \left(\frac{3}{4} \right) \text{ or allow } \pi \left(\frac{75}{4} + 50 \ln \left(\frac{3}{4} \right) \right)$	Applies limits of 4 and 1 and subtracts the correct way round. dM1 A1 oe [6] 10

Question 4 Notes	
4. (a)	<p>BE CAREFUL! Candidates will assign <i>their own</i> "A, B and C" for this question.</p> <p>B1 At least one of "B" or "C" are correct.</p> <p>B1 Breaks up their partial fraction correctly into three terms and both "B" = 25 and "C" = 100.</p> <p>Note If a candidate does not give partial fraction decomposition then: <ul style="list-style-type: none"> the 2nd B1 mark can follow from a correct identity. </p> <p>M1 Writes down a correct identity (although this can be implied) and attempts to find the value of either one of "A" or "B" or "C". This can be achieved by either substituting values into their identity or comparing coefficients and solving the resulting equations simultaneously.</p> <p>A1 Correct value for "A" which is found using a correct identity and follows from their partial fraction decomposition.</p> <p>Note If a candidate does not give partial fraction decomposition then the final A1 mark can be awarded for a correct "A" if a candidate writes out their partial fractions at the end.</p> <p>Note The correct partial fraction from no working scores B1B1M1A1.</p> <p>Note A number of candidates will start this problem by writing out the correct identity and then attempt to find "A" or "B" or "C". Therefore the B1 marks can be awarded from this method.</p> <p>Note Award SC B1B0M0A0 for $\frac{25}{x^2(2x+1)} \equiv \frac{B}{x^2} + \frac{C}{(2x+1)}$ leading to "B" = 25 or "C" = 100</p>
(b)	<p>B1 For a correct statement of $\pi \int \left(\frac{5}{x\sqrt{2x+1}} \right)^2$ or $\pi \int \frac{25}{x^2(2x+1)}$. Ignore limits and dx. Can be implied.</p> <p>Note The π can only be recovered later from a correct expression.</p>
	<p>For their partial fraction, (not $\sqrt{\text{their partial fraction}}$), where A, B, C are "their" part (a) constants</p> <p>M1 Either $\pm \frac{A}{x} \rightarrow \pm a \ln x$ or $\pm \frac{B}{x^2} \rightarrow \pm b x^{-1}$ or $\frac{C}{(2x+1)} \rightarrow \pm c \ln(2x+1)$.</p> <p>Note $\sqrt{\frac{B}{x^2}} \rightarrow \frac{\sqrt{B}}{x}$ which integrates to $\sqrt{B} \ln x$ is not worthy of M1.</p> <p>A1ft At least two terms from any of $\pm \frac{A}{x}$ or $\pm \frac{B}{x^2}$ or $\frac{C}{(2x+1)}$ correctly integrated. Can be un-simplified.</p> <p>A1ft All 3 terms from $\pm \frac{A}{x}$, $\pm \frac{B}{x^2}$ and $\frac{C}{(2x+1)}$ correctly integrated. Can be un-simplified.</p> <p>Note The 1st A1 and 2nd A1 marks in part (b) are both follow through accuracy marks.</p>
	<p>dM1 Dependent on the previous M mark. Applies limits of 4 and 1 and subtracts the correct way round.</p> <p>A1 Final correct exact answer in the form $a + b \ln c$. i.e. either $\frac{75}{4}\pi + 50\pi \ln\left(\frac{3}{4}\right)$ or $50\pi \ln\left(\frac{3}{4}\right) + \frac{75}{4}\pi$ or $50\pi \ln\left(\frac{9}{12}\right) + \frac{75}{4}\pi$ or $\frac{75}{4}\pi - 50\pi \ln\left(\frac{4}{3}\right)$ or $\frac{75}{4}\pi + 25\pi \ln\left(\frac{9}{16}\right)$ etc. Also allow $\pi \left(\frac{75}{4} + 50 \ln\left(\frac{3}{4}\right) \right)$ or equivalent.</p> <p>Note A candidate who achieves full marks in (a), but then mixes up the correct constants when writing their partial fraction can only achieve a maximum of B1M1A1A0M1A0 in part (b).</p> <p>Note The π in the volume formula is only required for the B1 mark and the final A1 mark.</p>

4. (b) Alternative method of integration

$$V = \pi \int_1^4 \left(\frac{5}{x\sqrt{2x+1}} \right)^2 dx$$

$$\int \frac{25}{x^2(2x+1)} dx ; u = \frac{1}{x} \Rightarrow \frac{du}{dx} = -\frac{1}{x^2}$$

$$= \int \frac{-25}{\left(\frac{2}{u}+1\right)} du = \int \frac{-25}{\left(\frac{2+u}{u}\right)} du = \int \frac{-25u}{(2+u)} du = -25 \int \frac{2+u-2}{(2+u)} du$$

$$= -25 \int 1 - \frac{2}{(2+u)} du = -25(u - 2\ln(2+u))$$

$$\left\{ \int_1^4 \frac{25}{x^2(2x+1)} dx = [-25u + 50\ln(2+u)]_1^4 \right\}$$

$$= \left(-\frac{25}{4} + 50\ln\left(\frac{9}{4}\right) \right) - (-25 + 50\ln 3)$$

$$= 50\ln\left(\frac{9}{4}\right) - 50\ln 3 - \frac{25}{4} + 25$$

$$= 50\ln\left(\frac{3}{4}\right) + \frac{75}{4}$$

$$\text{So, } V = \frac{75}{4}\pi + 50\pi\ln\left(\frac{3}{4}\right)$$

B1 For $\pi \int \left(\frac{5}{x\sqrt{2x+1}} \right)^2$
Ignore limits and dx. Can be implied.

M1 Achieves $\pm \alpha \pm \frac{\beta}{(k+u)}$ and integrates to give either $\pm \alpha u$ or $\pm \beta \ln(k+u)$

A1 **Dependent on the M mark.**
Either $-25u$ or $50\ln(2+u)$

A1 $-25(u - 2\ln(2+u))$

dM1 Applies limits of $\frac{1}{4}$ and 1 in u or 4 and 1 in x in their integrated function and subtracts the correct way round.

A1 $\frac{75}{4}\pi + 50\pi\ln\left(\frac{3}{4}\right)$ or allow $\pi\left(\frac{75}{4} + 50\ln\left(\frac{3}{4}\right)\right)$

Question Number	Scheme	Mark	
5. (a)	From question, $V = \frac{4}{3}\pi r^3$, $S = 4\pi r^2$, $\frac{dV}{dt} = 3$ $\left\{ V = \frac{4}{3}\pi r^3 \Rightarrow \right\} \frac{dV}{dr} = 4\pi r^2$ $\frac{dV}{dr} = 4\pi r^2$ (Can be implied)	B1 oe	
	$\left\{ \frac{dV}{dr} \times \frac{dr}{dt} = \frac{dV}{dt} \Rightarrow \right\} (4\pi r^2) \frac{dr}{dt} = 3$ $\left\{ \frac{dr}{dt} = \frac{dV}{dt} \div \frac{dV}{dr} \Rightarrow \right\} \frac{dr}{dt} = (3) \frac{1}{4\pi r^2}; \left\{ = \frac{3}{4\pi r^2} \right\}$ or $3 \div \text{Candidate's } \frac{dV}{dr}$;	M1 oe	
	When $r = 4\text{cm}$, $\frac{dr}{dt} = \frac{3}{4\pi(4)^2} \left\{ = \frac{3}{64\pi} \right\}$	dependent on previous M1. see notes	dM1
	Hence, $\frac{dr}{dt} = 0.01492077591\dots(\text{cm}^2 \text{s}^{-1})$	anything that rounds to 0.0149	A1 [4]
(b)	$\left\{ \frac{dS}{dt} = \frac{dS}{dr} \times \frac{dr}{dt} = \right\} \Rightarrow \frac{dS}{dt} = 8\pi r \times \frac{3}{4\pi r^2} \left\{ \text{or } \frac{6}{r} \text{ or } 8\pi r \times 0.0149\dots \right\}$ $8\pi r \times \text{Candidate's } \frac{dr}{dt}$	M1; oe	
	When $r = 4\text{cm}$, $\frac{dr}{dt} = 8\pi(4) \times \frac{3}{4\pi(4)^2}$ or $\frac{6}{4}$ or $8\pi(4) \times 0.0149\dots$		
	Hence, $\frac{dS}{dt} = 1.5 (\text{cm}^2 \text{s}^{-1})$	anything that rounds to 1.5	A1 cso [2]

6

Question 5 Notes

(a)	B1	$\frac{dV}{dr} = 4\pi r^2$ Can be implied by later working.
	M1	$\left(\text{Candidate's } \frac{dV}{dr} \right) \times \frac{dr}{dt} = 3$ or $3 \div \text{Candidate's } \frac{dV}{dr}$
	dM1	(dependent on the previous method mark) Substitutes $r = 4$ into an expression which is a result of a quotient of “3” and their $\frac{dV}{dr}$.
	A1	anything that rounds to 0.0149 (units are not required)
(b)	M1	$8\pi r \times \text{Candidate's } \frac{dr}{dt}$
	A1	anything that rounds to 1.5 (units are not required). Correct solution only.
	Note	Using $\frac{dr}{dt} = 0.0149$ gives $\frac{dS}{dt} = 1.4979\dots$ which is fine for A1.

Question Number	Scheme	Marks
<p>6.</p> <p>(a)</p> <p>(b)</p>	<p> $l_1: \mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}, \quad l_2: \mathbf{r} = \begin{pmatrix} 7 \\ 0 \\ 7 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ -5 \\ 4 \end{pmatrix} \quad \overline{OA} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \overline{OB} = \begin{pmatrix} 4 \\ p \\ 3 \end{pmatrix}$ </p> <p> $\{B \text{ lies on } l_2 \Rightarrow \mu = -1 \Rightarrow\} \quad p = 5$ </p> <p> $\{l_1 = l_2 \Rightarrow\} \begin{cases} \mathbf{i}: & 1 = 7 + 3\mu \\ \mathbf{j}: & 2 + 2\lambda = -5\mu \\ \mathbf{k}: & 3 - \lambda = 7 + 4\mu \end{cases}$ </p> <p>e.g. $\mathbf{i}: 7 + 3\mu = 1$</p> <p>So, $\mu = -2$</p> <p>Point of intersection is $\overline{OC} = \mathbf{i} + 10\mathbf{j} - \mathbf{k}$</p> <p>Finds $\lambda = 4$ and either</p> <ul style="list-style-type: none"> checks $\lambda = 4$ and $\mu = -2$ is true for the third component. substitutes $\mu = -2$ into l_1 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ and substitutes $\lambda = 4$ into l_2 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ 	<p>A lies on l_1 and B lies on l_2</p> <p>$p = 5$</p> <p>B1</p> <p>[1]</p> <p>Writes down an equation involving only one parameter. $\mu = -2$</p> <p>M1 A1 B1</p> <p>B1</p> <p>[4]</p>
<p>(b)</p>	<p>Alternative Method: Solving \mathbf{j} and \mathbf{k} simultaneously gives</p> <p>$8 = 14 + 3\mu$ or $23 + 3\lambda = 35$</p> <p>So, $\mu = -2$ or $\lambda = 4$</p> <p>Point of intersection is $\overline{OC} = \mathbf{i} + 10\mathbf{j} - \mathbf{k}$</p> <p>Finds $\lambda = 4$ and either</p> <ul style="list-style-type: none"> checks $\mu = -2$ is true for the \mathbf{i} component. substitutes $\mu = -2$ into l_1 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ and substitutes $\lambda = 4$ into l_2 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ 	<p>Writes down an equation involving only one parameter. Either $\mu = -2$ or $\lambda = 4$</p> <p>M1 A1 B1</p> <p>B1</p> <p>[4]</p>
<p>(c)</p> <p>(d)</p>	<p> $\overline{AC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix}$ and $\overline{BC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix} = \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix}$ </p> <p> $\pm \left(\begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix} \cdot \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix} \right)$ </p> <p> $\cos ACB = \frac{\overline{AC} \cdot \overline{BC}}{ \overline{AC} \cdot \overline{BC} } = \frac{0 + 40 + 16}{\sqrt{(0)^2 + (8)^2 + (-4)^2} \cdot \sqrt{(-3)^2 + (5)^2 + (-4)^2}}$ </p> <p> $\left\{ \cos ACB = \frac{0 + 40 + 16}{\sqrt{80} \cdot \sqrt{50}} = \frac{56}{\sqrt{4000}} \Rightarrow \right\} ACB = 27.69446... = 27.7$ (3 sf) </p> <p> $\text{Area } ACB = \frac{1}{2}(\sqrt{80})(\sqrt{50})\sin 27.69446...^\circ = 14.696888...$ </p>	<p>An attempt to find both the vectors (\overline{AC} or \overline{CA}) and (\overline{BC} or \overline{CB}).</p> <p>Applies dot product formula between their (\overline{AC} or \overline{CA}) and their (\overline{BC} or \overline{CB}).</p> <p>Anything that rounds to 27.7</p> <p>See notes Anything that rounds to 14.7</p> <p>M1 M1 A1</p> <p>A1</p> <p>[3]</p> <p>M1 A1</p> <p>[2] 10</p>

Question 6: Alternative Methods for Part (c)

<p>6. (c)</p>	<p>Alternative Method 1: Using the direction vectors of Line 1 and Line 2</p> $\mathbf{d}_1 = \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}, \quad \mathbf{d}_2 = \begin{pmatrix} 3 \\ -5 \\ 4 \end{pmatrix}$ $\cos \theta = \frac{\mathbf{d}_1 \cdot \mathbf{d}_2}{ \mathbf{d}_1 \mathbf{d}_2 } = \frac{\begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -5 \\ 4 \end{pmatrix}}{\sqrt{(0)^2 + (2)^2 + (-1)^2} \cdot \sqrt{(3)^2 + (-5)^2 + (4)^2}}$ $\left\{ \cos \theta = \frac{0 - 10 - 4}{\sqrt{5} \cdot \sqrt{50}} = \frac{-7\sqrt{10}}{25} \Rightarrow \right\} \theta = 152.3054385\dots$ <p>Angle $ACB = 180 - 152.3054385\dots = 27.69446145\dots = 27.7$ (3 sf)</p>	<p>Applies dot product formula between their \mathbf{d}_1 and \mathbf{d}_2</p> <p>M2</p> <p>Anything that rounds to 27.7</p> <p>A1</p> <p>[3]</p>
	<p>Alternative Method 2: The Cosine Rule</p> $\overrightarrow{AC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix} \quad \text{and} \quad \overrightarrow{BC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix} = \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix}$ <p>Also $\overrightarrow{AB} = \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 0 \end{pmatrix}$</p> <p>Note $\overrightarrow{AC} = \sqrt{80}$, $\overrightarrow{BC} = \sqrt{50}$ and $\overrightarrow{AB} = \sqrt{18}$</p> $(\sqrt{18})^2 = (\sqrt{80})^2 + (\sqrt{50})^2 - 2(\sqrt{80})(\sqrt{50})\cos \theta$ $\left\{ \cos \theta = \frac{7\sqrt{10}}{25} \right\} \Rightarrow \theta = 27.69446145\dots = 27.7$ (3 sf)	<p>An attempt to find both the vectors (\overrightarrow{AC} or \overrightarrow{CA}) and (\overrightarrow{BC} or \overrightarrow{CB}).</p> <p>M1</p> <p>Applies the cosine rule the correct way round.</p> <p>Anything that rounds to 27.7</p> <p>M1 oe</p> <p>A1</p> <p>[3]</p>
	<p>Alternative Method 3: Vector Cross Product</p> <p>Only apply this scheme if it is clear that a candidate is applying a vector cross product method.</p> $\overrightarrow{AC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix} \quad \text{and} \quad \overrightarrow{BC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix} = \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix}$ $\overrightarrow{AC} \times \overrightarrow{BC} = \begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix} \times \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix} = \left\{ \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 8 & -4 \\ -3 & 5 & -4 \end{vmatrix} = 24\mathbf{i} + 12\mathbf{j} + 24\mathbf{k} \right\}$ $\sin ACB = \frac{\sqrt{(24)^2 + (12)^2 + (12)^2}}{\sqrt{(0)^2 + (8)^2 + (-4)^2} \cdot \sqrt{(-3)^2 + (5)^2 + (-4)^2}}$ $\left\{ \sin ACB = \frac{\sqrt{864}}{\sqrt{80} \cdot \sqrt{50}} = \frac{3\sqrt{15}}{25} \Rightarrow \right\} \theta = 27.69446145\dots = 27.7$ (3 sf)	<p>An attempt to find both the vectors (\overrightarrow{AC} or \overrightarrow{CA}) and (\overrightarrow{BC} or \overrightarrow{CB}).</p> <p>M1</p> <p>Full method for applying the vector cross product formula between their (\overrightarrow{AC} or \overrightarrow{CA}) and their (\overrightarrow{BC} or \overrightarrow{CB}).</p> <p>M1</p> <p>Anything that rounds to 27.7</p> <p>A1</p> <p>[3]</p>

Question 6 Notes		
6. (a)	B1	$p = 5$ (Ignore working.)
(b)		Method 1
	M1	Writes down an equation involving only one parameter. This equation will usually be $7 + 3\mu = 1$ which is found from equating the i components of l_1 and l_2 .
	A1	Finds $\mu = -2$
	B1	Point of intersection of $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$. Allow $(1, 10, -1)$ or $\begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix}$.
	B1	Finds $\lambda = 4$ and either <ul style="list-style-type: none"> checks $\lambda = 4$ and $\mu = -2$ is true for the third component. substitutes $\mu = -2$ into l_1 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ and substitutes $\lambda = 4$ into l_2 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$
(b)		Alternative Method
	M1	Writes down an equation involving only one parameter. Solving the j and k components simultaneously will usually give either $8 = 14 + 3\mu$ or $23 + 3\lambda = 35$
	A1	Finds either $\mu = -2$ or $\lambda = 4$
	B1	Point of intersection of $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$. Allow $(1, 10, -1)$ or $\begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix}$.
	B1	Finds $\lambda = 4$ and either <ul style="list-style-type: none"> checks $\mu = -2$ is true for the i component. substitutes $\mu = -2$ into l_1 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ and substitutes $\lambda = 4$ into l_2 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$
(c)	M1	An attempt to find both the vectors $(\overline{AC}$ or $\overline{CA})$ and $(\overline{BC}$ or $\overline{CB})$ by subtracting.
	M1	Applies dot product formula between their $(\overline{AC}$ or $\overline{CA})$ and their $(\overline{BC}$ or $\overline{CB})$.
	A1	anything that rounds to 27.7
	Note	An answer of 0.48336... in radians without the correct answer in degrees is A0.
	Note	Some candidates will apply the dot product formula between vectors which are the wrong way round and achieve 152.3054385...°. If they give the acute equivalent of awrt 27.7 then award A1.
(d)	M1	$\frac{1}{2}(\text{their length } AC)(\text{their length } BC)\sin(\text{their } 27.7^\circ \text{ from part (c)})$
	A1	anything that rounds to 14.7. Also allow $6\sqrt{6}$.
	Note	Area $ACB = \frac{1}{2}(\sqrt{80})(\sqrt{50})\sin(152.3054385...^\circ) = \text{awrt } 14.7$ is M1A1.

Question Number	Scheme	Marks												
7.	$\frac{dN}{dt} = \frac{(kt-1)(5000-N)}{t}, \quad t > 0, \quad 0 < N < 5000$													
(a)	$\int \frac{1}{5000-N} dN = \int \frac{(kt-1)}{t} dt \quad \left\{ \text{or} = \int \left(k - \frac{1}{t} \right) dt \right\}$ <p>See notes</p> $-\ln(5000-N) = kt - \ln t; + c$ <p>See notes</p> <table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width:33%;"><i>then eg either...</i></td> <td style="width:33%;"><i>or...</i></td> <td style="width:33%;"><i>or...</i></td> </tr> <tr> <td>$-kt + c = \ln(5000-N) - \ln t$</td> <td>$kt + c = \ln t - \ln(5000-N)$</td> <td>$\ln(5000-N) = -kt + \ln t + c$</td> </tr> <tr> <td>$-kt + c = \ln\left(\frac{5000-N}{t}\right)$</td> <td>$kt + c = \ln\left(\frac{t}{5000-N}\right)$</td> <td>$5000-N = e^{-kt + \ln t + c}$</td> </tr> <tr> <td>$e^{-kt+c} = \frac{5000-N}{t}$</td> <td>$e^{kt+c} = \frac{t}{5000-N}$</td> <td>$5000-N = te^{-kt+c}$</td> </tr> </table> <p>leading to $N = 5000 - Ate^{-kt}$ with no incorrect working/statements. See notes</p>	<i>then eg either...</i>	<i>or...</i>	<i>or...</i>	$-kt + c = \ln(5000-N) - \ln t$	$kt + c = \ln t - \ln(5000-N)$	$\ln(5000-N) = -kt + \ln t + c$	$-kt + c = \ln\left(\frac{5000-N}{t}\right)$	$kt + c = \ln\left(\frac{t}{5000-N}\right)$	$5000-N = e^{-kt + \ln t + c}$	$e^{-kt+c} = \frac{5000-N}{t}$	$e^{kt+c} = \frac{t}{5000-N}$	$5000-N = te^{-kt+c}$	<p>B1</p> <p>M1 A1; A1</p> <p>A1 * cso</p> <p>[5]</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[4]</p>
<i>then eg either...</i>	<i>or...</i>	<i>or...</i>												
$-kt + c = \ln(5000-N) - \ln t$	$kt + c = \ln t - \ln(5000-N)$	$\ln(5000-N) = -kt + \ln t + c$												
$-kt + c = \ln\left(\frac{5000-N}{t}\right)$	$kt + c = \ln\left(\frac{t}{5000-N}\right)$	$5000-N = e^{-kt + \ln t + c}$												
$e^{-kt+c} = \frac{5000-N}{t}$	$e^{kt+c} = \frac{t}{5000-N}$	$5000-N = te^{-kt+c}$												
(b)	<p>$\{t = 1, N = 1200 \Rightarrow\} \quad 1200 = 5000 - Ae^{-k}$</p> <p>$\{t = 2, N = 1800 \Rightarrow\} \quad 1800 = 5000 - 2Ae^{-2k}$</p> <p>So $Ae^{-k} = 3800$</p> <p>and $2Ae^{-2k} = 3200$ or $Ae^{-2k} = 1600$</p> <p>Eg. $\frac{e^{-k}}{2e^{-2k}} = \frac{3800}{3200}$ or $\frac{2e^{-2k}}{e^{-k}} = \frac{3200}{3800}$</p> <p>So $\frac{1}{2}e^k = \frac{3800}{3200}$ or $2e^{-k} = \frac{3200}{3800}$</p> <p>$k = \ln\left(\frac{7600}{3200}\right)$ or equivalent $\left\{ \text{eg } k = \ln\left(\frac{19}{8}\right) \right\}$</p> <p>$\left\{ A = 3800(e^k) = 3800\left(\frac{19}{8}\right) \Rightarrow \right\} A = 9025$</p>	<p>At least one correct statement written down using the boundary conditions</p> <p>An attempt to eliminate A by producing an equation in only k.</p> <p>At least one of $A = 9025$ cao</p> <p>or $k = \ln\left(\frac{7600}{3200}\right)$ or exact equivalent</p> <p>Both $A = 9025$ cao</p> <p>or $k = \ln\left(\frac{7600}{3200}\right)$ or exact equivalent</p>												
(c)	<p>Alternative Method for the M1 mark in (b)</p> $e^{-k} = \frac{3800}{A}$ $2A\left(\frac{3800}{A}\right)^2 = 3200$ <p>An attempt to eliminate k by producing an equation in only A</p> <p>$\left\{ t = 5, N = 5000 - 9025(5)e^{-5\ln\left(\frac{19}{8}\right)} \right\}$</p> <p>$N = 4402.828401... = 4400$ (fish) (nearest 100)</p> <p>anything that rounds to 4400</p>	<p>M1</p> <p>B1</p> <p>[1]</p> <p>10</p>												

Question 7 Notes	
7. (a)	<p>B1 Separates variables as shown. dN and dt should be in the correct positions, though this mark can be implied by later working. Ignore the integral signs.</p> <p>M1 Either $\pm \lambda \ln(5000 - N)$ or $\pm \lambda \ln(N - 5000)$ or $kt - \ln t$ where $\lambda \neq 0$ is a constant.</p> <p>A1 For $-\ln(5000 - N) = kt - \ln t$ or $\ln(5000 - N) = -kt + \ln t$ or $-\frac{1}{k} \ln(5000 - N) = t - \frac{1}{k} \ln t$ oe</p> <p>A1 which is dependent on the 1st M1 mark being awarded.</p> <p>For applying a constant of integration, eg. $+c$ or $+\ln e^c$ or $+\ln c$ or A to their integrated equation $+c$ can be on either side of their equation for the 2nd A1 mark.</p> <p>Note</p> <p>A1 Uses a constant of integration eg. "c" or "$\ln e^c$" "$\ln c$" or and applies a fully correct method to prove the result $N = 5000 - Ate^{-kt}$ with no incorrect working seen. (Correct solution only.)</p> <p>NOTE IMPORTANT</p> <p>There needs to be an intermediate stage of justifying the A and the e^{-kt} in Ate^{-kt} by for example</p> <ul style="list-style-type: none"> • either $5000 - N = e^{\ln t - kt + c}$ • or $5000 - N = t e^{-kt + c}$ • or $5000 - N = t e^{-kt} e^c$ <p>or equivalent needs to be stated before achieving $N = 5000 - Ate^{-kt}$</p>
(b)	<p>B1 At least one of either $1200 = 5000 - Ae^{-k}$ (or equivalent) or $1800 = 5000 - 2Ae^{-2k}$ (or equivalent)</p> <p>M1</p> <ul style="list-style-type: none"> • Either an attempt to eliminate A by producing an equation in only k. • or an attempt to eliminate k by producing an equation in only A <p>A1 At least one of $A = 9025$ cao or $k = \ln\left(\frac{7600}{3200}\right)$ or equivalent</p> <p>A1 Both $A = 9025$ cao or $k = \ln\left(\frac{7600}{3200}\right)$ or equivalent</p> <p>Note Alternative correct values for k are $k = \ln\left(\frac{19}{8}\right)$ or $k = -\ln\left(\frac{8}{19}\right)$ or $k = \ln 7600 - \ln 3200$</p> <p>or $k = -\ln\left(\frac{3800}{9025}\right)$ or equivalent.</p> <p>Note $k = 0.8649...$ without a correct exact equivalent is A0.</p>
(c)	<p>B1 anything that rounds to 4400</p>

Question Number	Scheme	Marks
8.	$x = t - 4\sin t, \quad y = 1 - 2\cos t, \quad -\frac{2\pi}{3} \leq t \leq \frac{2\pi}{3}$ $A(k, 1)$ lies on the curve, $k > 0$	
(a)	$\left\{ \text{When } y=1, \right\} 1 = 1 - 2\cos t \Rightarrow t = -\frac{\pi}{2}, \frac{\pi}{2}$ $k \text{ (or } x) = \frac{\pi}{2} - 4\sin\left(\frac{\pi}{2}\right) \text{ or } x = -\frac{\pi}{2} - 4\sin\left(-\frac{\pi}{2}\right)$ $\left\{ \text{When } t = -\frac{\pi}{2}, k > 0, \right\}$ so $k = 4 - \frac{\pi}{2}$ or $\frac{8 - \pi}{2}$	<p>Sets $y=1$ to find t and uses their t to find x. M1</p> <p>x or $k = 4 - \frac{\pi}{2}$ A1</p>
(b)	$\frac{dx}{dt} = 1 - 4\cos t, \quad \frac{dy}{dt} = 2\sin t$	<p>At least one of $\frac{dx}{dt}$ or $\frac{dy}{dt}$ correct. B1</p> <p>Both $\frac{dx}{dt}$ and $\frac{dy}{dt}$ are correct. B1</p>
	<p>So, $\frac{dy}{dx} = \frac{2\sin t}{1 - 4\cos t}$</p> <p>At $t = -\frac{\pi}{2}, \frac{dy}{dx} = \frac{2\sin\left(-\frac{\pi}{2}\right)}{1 - 4\cos\left(-\frac{\pi}{2}\right)}; = -2$</p>	<p>Applies their $\frac{dy}{dt}$ divided by their $\frac{dx}{dt}$ and substitutes their t into their $\frac{dy}{dx}$. M1;</p> <p>Correct value for $\frac{dy}{dx}$ of -2 A1</p>
(c)	<p>$\frac{2\sin t}{1 - 4\cos t} = -\frac{1}{2}$</p> <p>gives $4\sin t - 4\cos t = -1$</p> <p>So $4\sqrt{2}\sin\left(t - \frac{\pi}{4}\right); = -1$ or $-4\sqrt{2}\cos\left(t + \frac{\pi}{4}\right); = -1$</p> <p>$t = \sin^{-1}\left(\frac{-1}{4\sqrt{2}}\right) + \frac{\pi}{4}$ or $t = \cos^{-1}\left(\frac{1}{4\sqrt{2}}\right) - \frac{\pi}{4}$</p> <p>$t = 0.6076875626... = 0.6077$ (4 dp)</p>	<p>Sets their $\frac{dy}{dx} = -\frac{1}{2}$ M1</p> <p>See notes A1</p> <p>See notes M1; A1</p> <p>See notes dM1</p> <p>anything that rounds to 0.6077 A1</p>
Question 8 Notes		
(c)	<p>NOTE <u>VERY IMPORTANT NOTE FOR PART (c)</u></p> <p>Candidates who state $t = 0.6077$ with no intermediate working from $4\sin t - 4\cos t = -1$ will get 2nd M0, 2nd A0, 3rd M0, 3rd A0.</p> <p>They will not express $4\sin t - 4\cos t$ as either $4\sqrt{2}\sin\left(t - \frac{\pi}{4}\right)$ or $-4\sqrt{2}\cos\left(t + \frac{\pi}{4}\right)$.</p> <p>OR use any acceptable alternative method to achieve $t = 0.6077$</p> <p>NOTE Alternative methods for part (c) are given on the next page.</p>	

[2]

[4]

[6]
12

Question 8: Alternative Methods for Part (c)	
8. (c)	<p>Alternative Method 1:</p> $\frac{2 \sin t}{1 - 4 \cos t} = -\frac{1}{2}$ <p style="text-align: right;">Sets their $\frac{dy}{dx} = -\frac{1}{2}$ M1</p> <p>eg. $\left(\frac{2 \sin t}{1 - 4 \cos t}\right)^2 = \frac{1}{4}$ or $(4 \sin t)^2 = (4 \cos t - 1)^2$ Squaring to give a correct equation. A1 This mark can be implied by a "squared" correct equation.</p> <p>or $(4 \sin t + 1)^2 = (4 \cos t)^2$ etc.</p> <p style="text-align: right;">Note: You can also give 1st A1 in this method for $4 \sin t - 4 \cos t = -1$ as in the main scheme.</p> <p style="text-align: center;">Squares their equation, applies $\sin^2 t + \cos^2 t = 1$ and achieves a three term quadratic equation of the form $\pm a \cos^2 t \pm b \cos t \pm c = 0$ M1 or $\pm a \sin^2 t \pm b \sin t \pm c = 0$ or eg. $\pm a \cos^2 t \pm b \cos t = \pm c$ where $a \neq 0, b \neq 0$ and $c \neq 0$.</p> <ul style="list-style-type: none"> • Either $32 \cos^2 t - 8 \cos t - 15 = 0$ • or $32 \sin^2 t + 8 \sin t - 15 = 0$ For a correct three term quadratic equation. A1 • Either $\cos t = \frac{8 \pm \sqrt{1984}}{64} = \frac{1 + \sqrt{31}}{8} \Rightarrow t = \cos^{-1}(\dots)$ which is dependent on the 2nd M1 mark. dM1 Uses correct algebraic processes to give $t = \dots$ • or $\sin t = \frac{-8 \pm \sqrt{1984}}{64} = \frac{-1 \pm \sqrt{31}}{8} \Rightarrow t = \sin^{-1}(\dots)$ anything that rounds to 0.6077 A1 $t = 0.6076875626\dots = 0.6077$ (4 dp) <p style="text-align: right;">[6]</p>
8. (c)	<p>Alternative Method 2:</p> $\frac{2 \sin t}{1 - 4 \cos t} = -\frac{1}{2}$ <p style="text-align: right;">Sets their $\frac{dy}{dx} = -\frac{1}{2}$ M1</p> <p>eg. $(4 \sin t - 4 \cos t)^2 = (-1)^2$ Squaring to give a correct equation. A1 This mark can be implied by a correct equation. Note: You can also give 1st A1 in this method for $4 \sin t - 4 \cos t = -1$ as in the main scheme.</p> <p>So $16 \sin^2 t - 32 \sin t \cos t + 16 \cos^2 t = 1$</p> <p style="text-align: right;">Squares their equation, applies both $\sin^2 t + \cos^2 t = 1$ and $\sin 2t = 2 \sin t \cos t$ and then achieves an equation of the form $\pm a \pm b \sin 2t = \pm c$ M1</p> <p>leading to $16 - 16 \sin 2t = 1$</p> <p style="text-align: right;">$16 - 16 \sin 2t = 1$ or equivalent. A1</p> <p>$\left\{ \sin 2t = \frac{15}{16} \Rightarrow \right\} t = \frac{\sin^{-1}(\dots)}{2}$ which is dependent on the 2nd M1 mark. dM1 Uses correct algebraic processes to give $t = \dots$</p> <p>$t = 0.6076875626\dots = 0.6077$ (4 dp) anything that rounds to 0.6077 A1</p> <p style="text-align: right;">[6]</p>

Question 8 Notes

8. (a)	M1	Sets $y = 1$ to find t and uses their t to find x .
	Note	M1 can be implied by either x or $k = 4 - \frac{\pi}{2}$ or 2.429... or $\frac{\pi}{2} - 4$ or $-2.429...$
	A1	x or $k = 4 - \frac{\pi}{2}$ or $\frac{8 - \pi}{2}$
	Note	A decimal answer of 2.429... (without a correct exact answer) is A0.
	Note	Allow A1 for a candidate using $t = \frac{\pi}{2}$ to find $x = \frac{\pi}{2} - 4$ and then stating that k must be $4 - \frac{\pi}{2}$ o.e.
(b)	B1	At least one of $\frac{dx}{dt}$ or $\frac{dy}{dt}$ correct. Note: that this mark can be implied from their working.
	B1	Both $\frac{dx}{dt}$ and $\frac{dy}{dt}$ are correct. Note: that this mark can be implied from their working.
	M1	Applies their $\frac{dy}{dt}$ divided by their $\frac{dx}{dt}$ and attempts to substitute their t into their expression for $\frac{dy}{dx}$.
	Note	This mark may be implied by their final answer. i.e. $\frac{dy}{dx} = \frac{2 \sin t}{1 - 4 \cos t}$ followed by an answer of -2 (from $t = -\frac{\pi}{2}$) or 2 (from $t = \frac{\pi}{2}$)
	Note	Applying $\frac{dx}{dt}$ divided by their $\frac{dy}{dt}$ is M0, even if they state $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$.
	A1	Using $t = -\frac{\pi}{2}$ (and not $t = \frac{3\pi}{2}$) to find a correct $\frac{dy}{dx}$ of -2 by correct solution only .
(c)	NOTE	If a candidate uses an incorrect $\frac{dy}{dx}$ expression in part (c) then the accuracy marks are not obtainable.
	1st M1	Sets their $\frac{dy}{dx} = -\frac{1}{2}$
	1st A1	Rearranges to give the correct equation with $\sin t$ and $\cos t$ on the same side . eg. $4 \sin t - 4 \cos t = -1$ or $4 \cos t - 4 \sin t = 1$ or $\sin t - \cos t = -\frac{1}{4}$ or $\cos t - \sin t = \frac{1}{4}$ or $4 \sin t - 4 \cos t + 1 = 0$ or $4 \cos t - 4 \sin t - 1 = 0$ or $\sin t - \cos t + \frac{1}{4} = 0$ etc. are fine for A1.
	2nd M1	Rewrites $\pm \lambda \sin t \pm \mu \cos t$ in the form of either $R \cos(t \pm \alpha)$ or $R \sin(t \pm \alpha)$ where $R \neq 1$ or 0 and $\alpha \neq 0$
	2nd A1	Correct equation. Eg. $4\sqrt{2} \sin\left(t - \frac{\pi}{4}\right) = -1$ or $-4\sqrt{2} \cos\left(t + \frac{\pi}{4}\right) = -1$ or $\sqrt{2} \sin\left(t - \frac{\pi}{4}\right) = -\frac{1}{4}$ or $\sqrt{2} \cos\left(t + \frac{\pi}{4}\right) = \frac{1}{4}$, etc.
	Note	Unless recovered, give A0 for $4\sqrt{2} \sin(t - 45^\circ) = -1$ or $-4\sqrt{2} \cos(t + 45^\circ) = -1$, etc.
	3rd M1	which is dependent on the 2nd M1 mark. Uses correct algebraic processes to give $t = \dots$
	4th A1	anything that rounds to 0.6077
	Note	Do not give the final A1 mark in (c) if there any extra solutions given in the range $-\frac{2\pi}{3} \leq t \leq \frac{2\pi}{3}$.
	Note	You can give the final A1 mark in (c) if extra solutions are given outside of $-\frac{2\pi}{3} \leq t \leq \frac{2\pi}{3}$.

