

Central limit theorem Mixed Exercise 5

1 By the central limit theorem $\bar{X} \approx \sim N\left(5, \frac{1}{100}\right)$, i.e. $\bar{X} \approx \sim N(5, 0.01)$

$$P(\bar{X} > 5.2) = 1 - P(\bar{X} < 5.2) \approx 1 - 0.9772 = 0.0228 \text{ (4 d.p.)}$$

$$2 \quad E(X) = \frac{1}{6}(1+2+4+5+7+8) = \frac{27}{6} = 4.5$$

$$\text{Var}(X) = \frac{1}{6}(1+2^2+4^2+5^2+7^2+8^2) - 4.5^2$$

$$= \frac{159}{6} - \frac{729}{36} = \frac{225}{36} = \frac{25}{4} = 6.25$$

By the central limit theorem $\bar{X} \approx \sim N\left(4.5, \frac{6.25}{20}\right)$, i.e. $\bar{X} \approx \sim N(4.5, 0.3125)$

$$P(\bar{X} < 4) \approx 0.1855 \text{ (4 d.p.)}$$

3 $X \sim N(1, 1)$ and by the central limit theorem $\bar{X} \sim N\left(1, \frac{1}{\sqrt{n}}\right)$

Standardise the sample mean.

$$P(\bar{X} < 0) = P\left(Z < -\sqrt{n}\right) \text{ and so require } P\left(Z > -\sqrt{n}\right) < 0.05$$

Using the table for the percentage points of the normal distribution:

$$P(Z = -1.645) = 0.05$$

$$\Rightarrow -\sqrt{n} < -1.645$$

$$\Rightarrow n > 2.706$$

So minimum sample size is $n = 3$ for the probability of a negative sample mean being less than 5%

4 Let the random variable X denote the number of sixes thrown by a student in 10 rolls of the dice, so

$$X \sim B\left(10, \frac{1}{6}\right)$$

$$E(X) = np = 10 \times \frac{1}{6} = \frac{5}{3}$$

$$\text{Var}(X) = np(1-p) = \frac{5}{3} \times \frac{5}{6} = \frac{25}{18}$$

By the central limit theorem $\bar{X} \approx \sim N\left(\frac{5}{3}, \frac{25}{18 \times 20}\right)$, i.e. $\bar{X} \approx \sim N\left(\frac{5}{3}, \frac{5}{72}\right)$

$$P(\bar{X} > 2) = 1 - P(\bar{X} < 2) \approx 1 - 0.8970 = 0.1030 \text{ (4 d.p.)}$$

5 a Let X be the number of buses that arrive in a 10-minute period, then $X \sim \text{Po}(2)$

$$P(X = 3) = \frac{e^{-2} 2^3}{3!} = 0.1804 \text{ (4 d.p.)}$$

5 b Let T be the number of buses that arrive in a two-hour period, so $T = 12\bar{X}$

By the central limit theorem $\bar{X} \approx \sim N\left(2, \frac{2}{12}\right)$, i.e. $\bar{X} \approx \sim N\left(2, \frac{1}{6}\right)$

$$P(T \geq 25) = P\left(\bar{X} \geq \frac{25}{12}\right)$$

$$P\left(\bar{X} \geq \frac{25}{12}\right) = 1 - P\left(\bar{X} < \frac{25}{12}\right) \approx 1 - 0.5809 = 0.4191 \text{ (4 d.p.)}$$

6 a Let the discrete random variable X be the number of children that a couple will have before having a daughter, then $X \sim \text{Geo}(0.5)$

$$P(X > 2) = (1 - 0.5)^2 = 0.5^2 = 0.25$$

$$\text{b } E(X) = \frac{1}{p} = \frac{1}{0.5} = 2$$

$$\text{Var}(X) = \frac{1-p}{p^2} = \frac{0.5}{0.5^2} = 2$$

By the central limit theorem $\bar{X} \approx \sim N\left(2, \frac{2}{10}\right)$, i.e. $\bar{X} \approx \sim N(2, 0.2)$

If $\bar{X} > 2.4$, the 10 couples will have more than 24 children

$$P(\bar{X} > 2.4) = 1 - P(\bar{X} < 2.4) \approx 1 - 0.8145 = 0.1855 \text{ (4 d.p.)}$$

7 a Let the random variable X be the mass of an egg, then $X \sim N(60, 25)$ and $\bar{X} \sim N\left(60, \frac{25}{48}\right)$

$$P(\bar{X} > 59) = 1 - P(\bar{X} < 59) = 1 - 0.0829 = 0.9171 \text{ (4 d.p.)}$$

b The answer in part **a** is not an estimate because the sample is taken from a population that is normally distributed.

c Let the random variable Y is the number of double yolk eggs in a crate of 48 eggs, so $Y \sim B(48, 0.1)$

$$E(Y) = np = 48 \times 0.1 = 4.8$$

$$\text{Var}(Y) = np(1-p) = 4.8 \times 0.9 = 4.32$$

By the central limit theorem $\bar{Y} \approx \sim N\left(4.8, \frac{4.32}{30}\right)$, i.e. $\bar{Y} \approx \sim N(4.8, 0.144)$

The probability that the sample of 30 crates will contain fewer than 150 double-yolk eggs is $P(\bar{Y} < 5)$ as $30 \times 5 = 150$

$$P(\bar{Y} < 5) \approx 0.7009 \text{ (4 d.p.)}$$

8 Consider a sample of 100 cups of coffee, so $\bar{S} \sim N(4.9, 0.0064)$. One pack of milk powder will be sufficient, if $100\bar{S} < 500$, i.e. $\bar{S} < 5$

$$P(\bar{S} < 5) = 0.8944 \text{ (4 d.p.)}$$

9 Let the random variable be X , so by the central limit theorem $\bar{X} \approx \sim N\left(40, \frac{9}{n}\right)$

Required to find minimum n such that $P(\bar{X} > 42) < 0.05$

Standardise the sample mean using $Z = \frac{\bar{X} - \mu}{\sigma}$, $\mu = 40$ and $\sigma = \frac{3}{\sqrt{n}}$

So for $\bar{X} = 42$, $Z = \frac{(42 - 40)\sqrt{n}}{3} = \frac{2\sqrt{n}}{3}$ and $P(\bar{X} > 42) = P\left(Z > \frac{2\sqrt{n}}{3}\right)$

Using the table for the percentage points of the normal distribution;

$$P(Z > 1.6449) = 0.05$$

$$\text{So } \frac{2\sqrt{n}}{3} > 1.6449$$

$$\Rightarrow \sqrt{n} > 2.46735$$

$$\Rightarrow n > 6.0878\dots$$

So $n = 7$ is the minimum sample size required for $P(\bar{X} > 42) < 0.05$

10 Let the random variable be X , so by the central limit theorem $\bar{X} \approx \sim N\left(35, \frac{9}{20}\right)$

$$P(\bar{X} > 37) = 1 - P(\bar{X} < 37) \approx 1 - 0.9986 = 0.0014 \text{ (4 d.p.)}$$

11 a The table describes the distribution of X

x	0	1
$P(X = x)$	0.4	0.6

$$E(X) = 0.6, \text{ Var}(X) = 0.6 - 0.6^2 = 0.24$$

b By the central limit theorem $\bar{X} \approx \sim N\left(0.6, \frac{0.24}{500}\right)$, i.e. $\bar{X} \approx \sim N(0.6, 0.00048)$

$$\begin{aligned} P(\bar{X} > 0.63) + P(\bar{X} < 0.57) &= 1 - P(\bar{X} < 0.63) + P(\bar{X} < 0.57) \\ &\approx 1 - 0.91454 + 0.08545 = 0.1709 \text{ (4 d.p.)} \end{aligned}$$

11 c Required to find minimum n such that $P(0.57 < \bar{X} < 0.63) > 0.95$

Standardise the sample mean using $Z = \frac{\bar{X} - \mu}{\sigma}$, $\mu = 0.6$ and $\sigma = \sqrt{\frac{0.24}{n}}$

So for $\bar{X} = 42$, $Z = \frac{(42 - 40)\sqrt{n}}{3} = \frac{2\sqrt{n}}{3}$ and $P(\bar{X} > 42) = P\left(Z > \frac{2\sqrt{n}}{3}\right)$

So require $P\left(-\frac{0.03\sqrt{n}}{\sqrt{0.24}} < Z < \frac{0.03\sqrt{n}}{\sqrt{0.24}}\right) > 0.95$

$\Rightarrow 1 - 2P\left(Z < -\frac{0.03\sqrt{n}}{\sqrt{0.24}}\right) > 0.95$ (by the symmetry of the normal distribution)

$\Rightarrow P\left(Z < -\frac{0.03\sqrt{n}}{\sqrt{0.24}}\right) < 0.025$

Using the table for the percentage points of the normal distribution

$$P(Z < -1.960) = 0.025$$

$$\Rightarrow -\frac{0.03\sqrt{n}}{\sqrt{0.24}} < -1.960$$

$$\Rightarrow \sqrt{n} > \frac{1.960 \times \sqrt{0.24}}{0.03} \Rightarrow \sqrt{n} > 32.0066\dots$$

$$\Rightarrow n > 1024.42\dots$$

So $n = 1025$

Challenge

$X_1 + \dots + X_n \sim N(n\mu, n\sigma^2)$ and so

$$\bar{X} = \frac{1}{n}(X_1 + \dots + X_n) \sim N\left(\frac{n\mu}{n}, \frac{n\sigma^2}{n^2}\right) = N\left(\mu, \frac{\sigma^2}{n}\right)$$